

# 56

## INFORMATION REPORT INFORMATION REPORT

CENTRAL INTELLIGENCE AGENCY

This material contains information affecting the National Defense of the United States within the meaning of the Espionage Laws, Title 18, U.S.C. Secs. 793 and 794, the transmission or revelation of which in any manner to an unauthorized person is prohibited by law.

C-O-N-F-I-D-E-N-T-I-A-L

50X1-HUM

COUNTRY	USSR	REPORT	
SUBJECT	Ultraviolet Radiation of the Sun and the Transition Layer Between Chromosphere and Corona	DATE DISTR.	16 May 61
		NO. PAGES	1
		REFERENCES	
DATE OF INFO.			50X1-HUM
PLACE & DATE ACQ.			50X1-HUM

THIS IS UNEVALUATED INFORMATION

abstract of a paper entitled  
"Ultraviolet Radiation of the Sun and the Transition Layer Between  
Chromosphere and Corona" by GS Ivanov-Kholodny and GM Nikolskiy.

50X1-HUM

OFFICIAL USE ONLY.

- end -

C-O-N-F-I-D-E-N-T-I-A-L

STATE	ARMY	NAVY	AIR	FBI	AEC						
-------	------	------	-----	-----	-----	--	--	--	--	--	--

## INFORMATION REPORT INFORMATION REPORT

NOFORN

NO DISSEM ABROAD

LIMITED

Dissemination limited to full-time employees of CIA, AEC and FBI, and, within State and Defense, to the intelligence components, other offices producing NIS elements, and higher echelons with their immediate supporting staffs. Not to be disseminated to consultants, external projects or reserve personnel on short term active duty (excepting individuals who are normally full-time employees of CIA, AEC, FBI, State or Defense) unless the

50X1-HUM

Sanitized Copy Approved for Release 2011/02/28 : CIA-RDP80T00246A013000370001-6

**Page Denied**

Sanitized Copy Approved for Release 2011/02/28 : CIA-RDP80T00246A013000370001-6

# FOR OFFICIAL USE ONLY

## ULTRAVIOLET RADIATION OF THE SUN AND THE TRANSITION LAYER BETWEEN CHROMOSPHERE AND CORONA

by G.S.Ivanov-Kholodny and G.M.Nikolsky

### Abstract

1. Most of the solar ultraviolet lines are emitted by the transition region and the corona. In the spectral ranges of  $1000-2000 \text{ \AA}^\circ$  /1-6/ and  $84-1200 \text{ \AA}^\circ$  are recorded  $\sim 200$  lines by means of the rocket spectrographs /7/.

2. The calibration was carried out in absolute units and as a result of this there were obtained  $J(L_\alpha) = 3$ ,  $J(H_\epsilon \text{ II } \lambda 304) = 1.2$  and  $J(H_\epsilon \text{ II } \lambda 584) = 0.1 \text{ erg cm}^2 \text{ sec}^{-1}$ . These values are consistent with the recent ionospheric data according which the intensity of the ultraviolet radiation for  $\lambda < 900 \text{ \AA}^\circ$  is  $\sim 30 \text{ erg cm}^2 \text{ sec}^{-1}$  /8/.

3. From the theoretical consideration of atomic ionisation and excitation in the transition region there follows an expression for the "luminosity function" of  $\Delta\varphi_i$  for a given ion:

$$\Delta\varphi_i = \left\{ n_e^2 T_i^{-3/2} (h_2 - h_1) \right\} = \frac{2.3 \cdot 10^{12} J \cdot \lambda}{\mathcal{X} \cdot W' \cdot f_{12}}, \quad (I)$$

where  $h_2$  and  $h_1$  are the upper and the lower heights of emitted layer accordingly,  $f_{12}$  - the oscillator strengths of the corresponding atomic transition,  $\lambda$  - wave length of an ultraviolet line and  $\mathcal{X}$  - the abundance of the element relative to the hydrogen. The probability of excitation of a given atomic level as an effect of electron collision is a function of  $\mathcal{X}$  or  $T$

$$W' = 10 \cdot T^{-3/2} \left\{ \frac{e^{-x}}{x} - E_i(x) \right\}, \quad (2)$$

where  $E_i(x) = \int_x^\infty \frac{e^{-x}}{x} dx$ ,  $x = \frac{h\nu}{kT}$ .

FOR OFFICIAL USE ONLY

- 2 -

4. Data regarding the 17 ultraviolet lines were used and two empirical relations were established

$$\Delta\varphi_i = f(T_i) \text{ (fig.1) and } T(h_2) - T(h_1) = 2 \cdot T_i^{0.92} \quad (3)$$

From fig.1 it is seen that the abundance of nitrogen  $\mathcal{Z}(N) \approx 10^{-5}$ , i.e. it is about 30 times less as compared to the one usually assumed.

5. The self absorption in the ultraviolet lines was found to be negligible although the optical depth may be large.

6. From this there were calculated ultraviolet and X-ray continuums, emitted by the transition region and the corona. It was concluded that according to the rocket observations the solar radiation in the spectral range of 30-1500 Å° mainly consists of line emission.

7. Radioemission as calculated using (4) is in accordance with the observations if one assumes, that during the maximum solar cycle at least 90% of ultraviolet radiation are emitted by active regions in case they occupy  $\sim 0.1$  of the solar surface and are  $\sim 100$  times brighter than the undisturbed region.

8. Eclipse data of Athay and al. / 9 /, were used for estimating a function:

$$\psi(h) = \int_h^{\infty} n_e^2 T^{-3/2} dh \quad (5)$$

for active and undisturbed regions of the chromosphere. From (4) and (5) the distributions of  $T$  and  $n_e$  with height were found, and thus models of active and undisturbed transition regions were constructed (fig.2). Temperature in the transition region changes with height smoother than in previous models.

(Preliminary results are published in /10/).

FOR OFFICIAL USE ONLY

R e f e r e n c e s :


**FOR OFFICIAL USE ONLY**

R e f e r e n c e s

Л и т е р а т у р а

- ~~W. Petri, Forschungsber. Obs., Wendelstein, Bd. 2, 15, 16, 1952~~
1. F.S. Johnson, J.D. Purcell, R. Tousey, Bull. Amer. Phys. Soc., 29, 33, 1954; Phys. Rev., 95, 621, 1954.
  2. F.S. Johnson, H.H. Malitson, J.D. Purcell, R. Tousey, Astrophys. J., 127, 80, 1958.
  3. A.S. Jursa, F.J. Le Blanc, Y. Tanaka, J. Opt. Soc. Amer., 45, 1085, 1955.
  4. W.E. Behring, H. McAlister, W.A. Rense, Astrophys. J., 127, 676, 1958.
  5. A. Aboud, W.E. Behring, W.A. Rense, Astrophys. J., 130, 381, 1959.
  6. H. Friedman, Sci. American, 200, 52, 1959.
  7. T. Violet, W.A. Rense, Astrophys. J., 130, 954, 1959.
  8. Г.С. Иванов-Холодный, Докл. АН СССР, 137, №2, 1961 (в печати)
  9. R.G. Aghay, D.H. Menzel, J.C. Pecker, R.N. Thomas, Astrophys. J. Suppl.,
  10. Г.С. Иванов-Холодный, Г.М. Никольский, Астрон. ж., 38, №1, <sup>(45)</sup>1961.

**FOR OFFICIAL USE ONLY**



FOR OFFICIAL USE ONLY

